

# Molecular Mechanisms of Cellular Adhesion in Tissue Morphogenesis

**Pat Wilson**

PhD

University of São Paulo  
São Paulo, 05508-070, Brazil

**Cameron Robinson**

Dr.

Indian Institute of Science  
Bangalore, Karnataka 560012, India

**Jesse Garcia**

Prof.

University of Cape Town  
Cape Town, 7700, South Africa

**Abstract.** Cellular adhesion is crucial for tissue morphogenesis, influencing cell positioning and tissue architecture. This article delves into the molecular mechanisms that govern cellular adhesion during tissue development. We examine the roles of adhesion molecules and their interactions with the extracellular matrix. Our study highlights the significance of cellular adhesion in morphogenesis and its implications for developmental biology.

**Keywords:** Cellular adhesion, Tissue morphogenesis, Adhesion molecules, Extracellular matrix, Developmental biology

## Introduction

Cellular adhesion is a fundamental biological process that enables cells to attach to each other and to the extracellular matrix, thereby playing a crucial role in tissue morphogenesis. The formation and maintenance of tissue architecture depend on precise cell-cell and cell-matrix interactions, mediated by a wide array of adhesion molecules. These molecules not only ensure structural integrity but also influence cell signaling pathways that regulate cell proliferation, differentiation, and migration. In this article, we explore the molecular mechanisms underlying cellular adhesion, focusing on the roles of key adhesion molecules and their interactions with the extracellular matrix. We discuss how these mechanisms contribute to tissue development and morphogenesis, as well as their implications for understanding developmental disorders. Our findings underscore the importance of cellular adhesion in developmental biology.

**This is a preliminary version. To read the full version of the article, please purchase a subscription.**

## References

1. Sharma, V. K., Basu, S., & Chakraborty, S. (2015). RNAi mediated broad-spectrum transgenic resistance in *Nicotiana benthamiana* to chilli-infecting begomoviruses. *Plant cell reports*, 34(8), 1389-1399.
2. Sharma, V. K., Kushwaha, N., Basu, S., Singh, A. K., & Chakraborty, S. (2015). Identification of siRNA generating hot spots in multiple viral suppressors to generate broad-spectrum antiviral resistance in plants. *Physiology and Molecular Biology of Plants*, 21(1), 9-18.
3. Kumar, R. V., Sharma, V. K., Chattopadhyay, B., & Chakraborty, S. (2012). An improved plant regeneration and *Agrobacterium*-mediated transformation of red pepper (*Capsicum annuum* L.). *Physiology and Molecular Biology of Plants*, 18(4), 357-364.
4. Sharma, V., Zheng, W., Huang, J., & Cook, D. E. (2020). CRISPR-Cas RNA targeting using transient Cas13a expression in *Nicotiana benthamiana*. In *RNA Abundance Analysis: Methods and Protocols* (pp. 1-18). New York, NY: Springer US.